
Granite Island in Lake Superior A new BSRN water site proposal

Bryan Fabbri¹, Fred Denn¹ and Greg Schuster² 1- Science Systems and Applications, Inc. (SSAI), 2- Nasa Langley Research Center 15th BSRN Scientific Review and Workshop – Boulder, Colorado, USA 2018 July 16-20



-160°

80°

60°

40°

20°

0°

-20°

-40°

-60°

-80°


-160°

Stations

* Running

Inactive

Closed Candidate



PSU

1 H *

BER

LRC 🏠

-80°

-60°

-40°

-20

80°

70°

60°


50°


40°

30°

20°

TO EN





Closest land point is about 10 km (6 miles) away.




Granite Island Coordinates: 46.721 N 87.411 W north

Granite Island information:

- 0.01 square km (2.5 acre) granite rock island
- Granite Island at its base is approximately 193 m above sea level
- Rises nearly perpendicular to 18 m above surface of Lake Superior
- Surrounded by deep water (18-30 m)



Transportation to Granite Island is by boat. Pictured is a 9 m Zodiac 920.






Installation as viewed from the boat.

Installation is approximately 15 m above the water line (as measured from the solar tracker height). The Granite Island installation is approximately 208 m above sea level.



Motivation for establishing a Measurement Site at Granite Island




NASA Langley is interested in using BSRN and Clouds and the Earth's Radiant Energy System (CERES) measurements with the Great Lakes Evaporation Network (GLEN) data to improve understanding of the Earth's energy budget.

Other motivations:

- Water sites are uncommon
- Surface validation of satellites such as CERES

CERES sample footprints



=

Sign in







Instruments Currently Collecting Data at Granite Island:


- Downwelling Shortwave Direct (K&Z CHP 1)
- Downwelling Shortwave Diffuse (K&Z CM 22)
- Downwelling Shortwave Global (K&Z CM 22)
- Downwelling Longwave (K&Z CG4)
- Solar Tracker (K&Z Solys2)
- AERONET cimel sunphotometer

Upcoming Instruments Planned:

• Meteorological (Temperature, RH, Pressure, Wind Speed and Direction)

Solar Power System at Granite Island







A week of downwelling shortwave and longwave data.

A single day of downwelling shortwave (direct, diffuse and global) and downwelling longwave.

?





A single day of AERONET data at Granite Island.




Closest upper-air sonde launch station (as the crow flies) to Granite Island is KGRB (Green Bay, Wisconsin) at 240 km.



JUN.22,18 03:06:00PM





Seagulls are an issue that will need to be resolved.



A temporary solution is using tie wraps to deter seagulls from perching on the domes.

Sun rise/set azimuths





Calibration Procedures

- Two Automatic Hickey-Frieden (AHF) cavity radiometers are used to calibrate the shortwave instruments every year at the Langley Research Center location (BSRN site ID: LRC).
- Have attended the National Pyrheliometer Comparison at the National Renewable Energy Laboratory (NREL) in Golden, Colorado every year to calibrate the AHF's for the last 20 years.
- Have attended the International Pyrheliometer Comparison in Davos, Switzerland every 5th year (last was in 2015) since 2000.
- The longwave instruments are sent to NREL for calibration every other year.

Acknowledgements:

- We would like to thank Scott Holman, owner of Granite Island for allowing a new measurement site to be introduced.
- We thank NASA Langley and the CERES project for providing the funding for Granite Island.
- We appreciate Brent Holben (AERONET PI) for providing an AERONET cimel at Granite Island.
- We thank John Lenters, Warren Smith, students and faculty of Northern Michigan University and especially Dan "Ducky" Chiconsky for assisting in the installation of instruments at Granite Island.



THANK YOU

Bryan Fabbri 1-757-951-1639 bryan.e.fabbri@nasa.gov

Fred Denn 1-757-951-1636 frederick.m.denn@nasa.gov

